

Storage: Store at room temperature

*Shelf Life: 3 years

Kampo product

Approval No.	16100AMZ04739000
Date of Initial Marketing in Japan	October 1987

N21

Kotaro Shohangekabukuryoto Extract Fine Granules

3. COMPOSITION AND PRODUCT DESCRIPTION

3.1 Composition

Brand name	Kotaro Shohangekabukuryoto Extract Fine Granules
Active ingredient	6.0g of Kotaro Shohangekabukuryoto Extract Fine Granules contains 1.2g of the dried extract of the following mixed crude drugs. JP Pinellia Tuber.....5.0g JP Ginger.....1.3g JP Poria Sclerotium.....5.0g (JP: Japanese Pharmacopoeia)
Excipients	Magnesium Stearate, Corn Starch, Lactose Hydrate, Pullulan, Magnesium Aluminometasilicate

3.2 Product Description

Dosage form	Fine granules
Color	Yellowish brown to milky white
Taste	Slightly sweet, then bitter
Odor	Characteristic odor
ID code	N21

4. INDICATIONS

Vomiting due to the specified symptoms and signs involving the digestive system and abdomen.

Mild hyperemesis gravidarum and emesis.

6. DOSAGE AND ADMINISTRATION

The usual adult dosage for oral use is 6.0g daily in 2 or 3 divided doses before or between meals.

The dosage may be adjusted according to the patient's age, body weight, and symptoms.

8. IMPORTANT PRECAUTIONS

8.1 When this product is used, the patient's "SHO"

(constitution/symptoms) should be taken into consideration. The patient's progress should be carefully monitored, and if no improvement in symptoms or findings is observed, continuous administration should be avoided.

8.2 When this product is used in combination with other Kampo products, etc., attention should be paid to the duplication of the contained crude drugs.

9. PRECAUTIONS CONCERNING PATIENTS WITH SPECIFIC BACKGROUNDS

9.5 Pregnant Women

This product should be used in pregnant women or women who may possibly be pregnant only if the expected therapeutic benefits outweigh the possible risks associated with treatment.

9.6 Breast-feeding Women

Considering the therapeutic benefits and the benefits of breastfeeding, continuation or discontinuation of breastfeeding should be considered.

9.7 Pediatric Use

No clinical studies have been conducted in children.

9.8 Geriatric Use

Since the physiological functions are generally decrease in elderly patients, careful supervision is recommended; measures such as reducing the dose may be considered.

20. PRECAUTIONS FOR HANDLING

- 20.1 To maintain the quality of the product, avoid moisture as much as possible and store in a cool place, away from direct sunlight.
- 20.2 Avoid moisture, especially after opening, and handle with care.
- 20.3 Since this product is made from crude drugs, the color of the product may vary.

22. PACKAGING

- 500g [bottle, loose]
2.0g×42 packets [sachets]
2.0g×231 packets [sachets]

24. REFERENCE REQUEST AND CONTACT INFORMATION

Pharmaceutical Division, Kotaro Pharmaceutical Co., Ltd.
5-23, Nakatsu 2-chome, Kita-ku, Osaka 531-0071, Japan

26. MARKETING AUTHORIZATION HOLDER, etc.

26.1 Marketing Authorization Holder

Kotaro Pharmaceutical Co., Ltd.
5-23, Nakatsu 2-chome, Kita-ku, Osaka 531-0071, Japan